



A fifth order block methods for solving second-order stiff ordinary differential equations using trigonometric functions and polynomial function as the basis function

Opeyemi O. Enoch , Catherine O. Alakofa *

Department of Mathematics, Federal University Oye-Ekiti, Ekiti State, Nigeria

Abstract

The numerical solution of second-order ordinary differential equations (ODEs) is examined in this work through a four-step linear multistep method. It employs a combination of trigonometric and polynomial functions as the approximate solution to the general second-order ordinary differential equations (ODEs). The method was developed using interpolation and collocation techniques. This methodology involves interpolating the basis function at two points and subsequently collocating it across all points, ensuring a robust scheme. To assess its efficacy, we solved three initial value problems (IVPs) associated with stiff differential equations. Through this examination, we established the method's core characteristics: consistency, zero stability, and consequently convergence. This thorough analysis demonstrates its reliability and suitability for resolving second-order ordinary differential equations. The comparison of our newly derived block method against existing approaches reveals its superiority. Our method's performance, evaluated across a spectrum of stiff second-order ordinary differential equations, surpasses the outcomes obtained from established authors. This substantiates its efficiency and effectiveness in addressing these mathematical challenges. This study marks a significant advancement by introducing a robust approach that not only accurately solves second-order ordinary differential equations but also streamlines the computational process. By integrating trigonometric and polynomial functions and leveraging interpolation and collocation techniques, our method stands out for its accuracy, stability, and convergence properties, offering a promising avenue for future research in this domain.

DOI:10.46481/asr.2024.3.2.156

Keywords: Trigonometric function, Polynomial function, Collocation, Interpolation, Stiff ODEs, Block method

Article History:

Received: 08 November 2023

Received in revised form: 29 February 2024

Accepted for publication: 29 May 2024

Published: 29 June 2024

© 2024 The Author(s). Published by the [Nigerian Society of Physical Sciences](#) under the terms of the [Creative Commons Attribution 4.0 International license](#). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

1. Introduction

Presented in this paper is the numerical solution of general second-order ordinary differential equations initial value problems (IVP) of the form:

$$y'' = f(x, y, y'), \quad y(a) = \eta_0, \quad y'(a) = \eta_1, \quad x \in [a, b]. \quad (1)$$

These second-order ODEs, which are used to solve numerous problems in physical sciences and engineering, are used to model a variety of natural processes. Many forms of the equation (1) may be difficult to solve analytically. As a result, numerical systems

*Corresponding author: Tel.: +234-806-0674-204.

Email address: alakofaoluwasseyi@gmail.com (Catherine O. Alakofa)

are frequently developed to solve them. Because a lot of problems of the form (1) are difficult to solve, approximate numerical integrations are commonly utilized to solve them. These second-order ODEs are frequently transformed to similar first-order ordinary differential equations and solved using the appropriate method. Among others who have discussed the reduction technique are Refs. [1–3].

To circumvent the problem of changing equation (1) to an equal system of first order ODEs, scholars like Awoyemi & Olanegan *et al.* [3, 4] proposed a linear multistep technique for solving equation (1) directly. According to Ref. [3], the continuous linear multistep method outperforms the discrete method in terms of error estimation by providing a simplified coefficient for more analytical work at various points and ensuring easy appropriation of solution at all interior points of the integration interval.

Authors such as Omar & Kuboye [6] and Onumanyi *et al.* [7] have presented continuous linear multistep techniques. These scholars used the predictor-corrector and Taylor series expansion to acquire starting values for their approaches. The predictor-corrector approach is expensive, according to Ref. [8], since subroutines are difficult to construct due to the specific methods necessary to offer beginning values and adjust the step size, resulting in longer computer time and more human effort. The correctors are not in the same order as the predictors. As a result, the accuracy of the method is very poor.

To rise above the drawbacks of the reduction and predictor-corrector methods, researchers such as Refs. [8–10] developed the block technique. The aim of this paper is to develop and implement a new continuous four step linear multistep block method that is zero-stable, consistent, and convergent for direct and accurate solution of second-order ODEs of initial value problems using trigonometric and polynomial functions as the basis function.

2. Derivation of the scheme

In deriving this method, basis function of the form,

$$y(x) = \sum_{n=0}^4 a_n x^n + a_5 \sin(\omega x) + a_6 \cos(\omega x), \quad (2)$$

is considered as the approximate solution to equation (1). The second derivative of equation (2) is obtained as:

$$y''(x) = \sum_{n=2}^4 n(n-1)a_n x^{n-2} - a_5(\omega^2) \sin(\omega x) - a_6(\omega^2) \cos(\omega x). \quad (3)$$

Collocating equation (3) at $x = x_{n+j}$, $j = 0(1)4$ and interpolating equation (2) at $x = x_{n+j}$, $j = 0, 1$. These equations are then combined together to give a non linear system of equations of the form:

$$\begin{bmatrix} 1 & x_n & x_n^2 & x_n^3 & x_n^4 & \sin wx_n & \cos wx_n \\ 1 & x_{n+1} & x_{n+1}^2 & x_{n+1}^3 & x_{n+1}^4 & \sin wx_{n+1} & \cos wx_{n+1} \\ 0 & 0 & 2 & 6x_n & 12x_n^2 & -w^2 \sin wx_n & -w^2 \cos wx_n \\ 0 & 0 & 2 & 6x_{n+1} & 12x_{n+1}^2 & -w^2 \sin wx_{n+1} & -w^2 \cos wx_{n+1} \\ 0 & 0 & 2 & 6x_{n+2} & 12x_{n+2}^2 & -w^2 \sin wx_{n+2} & -w^2 \cos wx_{n+2} \\ 0 & 0 & 2 & 6x_{n+3} & 12x_{n+3}^2 & -w^2 \sin wx_{n+3} & -w^2 \cos wx_{n+3} \\ 0 & 0 & 2 & 6x_{n+4} & 12x_{n+4}^2 & -w^2 \sin wx_{n+4} & -w^2 \cos wx_{n+4} \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \\ a_4 \\ a_5 \\ a_6 \end{bmatrix} = \begin{bmatrix} y_n \\ y_{n+1} \\ f_n \\ f_{n+1} \\ f_{n+2} \\ f_{n+3} \\ f_{n+4} \end{bmatrix}. \quad (4)$$

Gaussian elimination technique is used in finding the values of a'_n s in equation (4) which are then substituted into equation (2) to produce a continuous implicit method of the form

$$y(x) = \sum_{j=0}^k \alpha_j(x) y_{n+j} + h^2 \left[\sum_{j=0}^k \beta_j(x) f_{n+j} \right], \quad (5)$$

where the coefficients of y_{n+j} and f_{n+j} gives

$$\begin{aligned}
\alpha_0 &= -1, \\
\alpha_1 &= 2, \\
\beta_0 &= \frac{14 \sin(wh)h^2w^2 - 13h^2w^2 \sin(2wh) - 36 \sin(wh) + 36 \sin(2wh) - 12 \sin(3wh)}{60 \sin(wh)w^2 + 12w^2 \sin(3wh) - 48w^2 \sin(2wh)}, \\
\beta_1 &= \frac{33 \sin(wh)h^2w^2 - 12h^2w^2 \sin(2wh) + 13w^2h^2 \sin(3wh) + 84 \sin(wh) - 96 \sin(2wh) + 36 \sin(3wh)}{60 \sin(wh)w^2 + 12w^2 \sin(3wh) - 48w^2 \sin(2wh)}, \\
\beta_2 &= \frac{\sin(wh)h^2w^2 - w^2h^2 \sin(3wh) - 17h^2w^2 \sin(2wh) - 18 \sin(3wh) - 18 \sin(wh) + 36 \sin(2wh)}{30 \sin(wh)w^2 - 24w^2 \sin(2wh) + 6w^2 \sin(3wh)}, \\
\beta_3 &= \frac{21 \sin(wh)h^2w^2 + 12h^2w^2 \sin(2wh) + w^2h^2 \sin(3wh) - 36 \sin(wh) + 12 \sin(3wh)}{60 \sin(wh)w^2 + 12w^2 \sin(3wh) - 48w^2 \sin(2wh)}, \\
\beta_4 &= \frac{-10 \sin(wh)h^2w^2 - h^2w^2 \sin(2wh) + 24 \sin(wh) - 12 \sin(2wh)}{20 \sin(wh)w^2 + 4w^2 \sin(3wh) - 16w^2 \sin(2wh)}. \tag{6}
\end{aligned}$$

Letting $u = \omega h$, the converted coefficients of the method in series form is given as

$$\begin{aligned}
\beta_0 &= \frac{19}{240}h^2 + \frac{221}{60480}h^2u^2 + \frac{233}{1814400}h^2u^4 + \frac{199}{53222400}h^2u^6 + O(u^8), \\
\beta_1 &= \frac{17}{20}h^2 - \frac{79}{7560}h^2u^2 - \frac{79}{226800}h^2u^4 - \frac{61}{6652800}h^2u^6 + O(u^8), \\
\beta_2 &= \frac{7}{120}h^2 + \frac{19}{2016}h^2u^2 + \frac{83}{302400}h^2u^4 + \frac{1}{197120}h^2u^6 + O(u^8), \\
\beta_3 &= \frac{1}{60}h^2 - \frac{2}{945}h^2u^2 - \frac{1}{56700}h^2u^4 + \frac{1}{415800}h^2u^6 + O(u^8), \\
\beta_4 &= -\frac{1}{240}h^2 - \frac{31}{60480}h^2u^2 - \frac{67}{1814400}h^2u^4 - \frac{109}{53222400}h^2u^6 + O(u^8), \tag{7}
\end{aligned}$$

or y_{n+3} ,

$$\begin{aligned}
\alpha_0 &= -2, \\
\alpha_1 &= 3, \\
\beta_0 &= \frac{9h^2w^2 \sin(2wh) - 6 \sin(wh)h^2w^2 - 20 \sin(2wh) + 16 \sin(wh) + 8 \sin(3wh)}{16w^2 \sin(2wh) - 20 \sin(wh)w^2 - 4w^2 \sin(3wh)}, \\
\beta_1 &= \frac{8h^2w^2 \sin(2wh) - 37 \sin(wh)h^2w^2 - 9w^2h^2 \sin(3wh) + 48 \sin(2wh) - 24 \sin(wh) - 24 \sin(3wh)}{16w^2 \sin(2wh) - 20 \sin(wh)w^2 - 4w^2 \sin(3wh)}, \\
\beta_2 &= \frac{\sin(wh)h^2w^2 - w^2h^2 \sin(3wh) + 19h^2w^2 \sin(2wh) - 12 \sin(2wh) + 12 \sin(3wh) - 12 \sin(wh)}{8w^2 \sin(2wh) - 10 \sin(wh)w^2 - 2w^2 \sin(3wh)}, \\
\beta_3 &= \frac{29 \sin(wh)h^2w^2 + w^2h^2 \sin(3wh) + 8h^2w^2 \sin(2wh) - 56 \sin(wh) + 8 \sin(3wh) + 16 \sin(2wh)}{20 \sin(wh)w^2 + 4w^2 \sin(3wh) - 16w^2 \sin(2wh)}, \\
\beta_4 &= \frac{-10 \sin(wh)h^2w^2 - h^2w^2 \sin(2wh) + 24 \sin(wh) - 12 \sin(2wh)}{60 \sin(wh)w^2 + 12w^2 \sin(3wh) - 48w^2 \sin(2wh)}. \tag{8}
\end{aligned}$$

The converted coefficients of the method in series form is given as

$$\begin{aligned}
\beta_0 &= \frac{37}{240}h^2 + \frac{137}{20160}h^2u^2 + \frac{19}{86400}h^2u^4 + \frac{289}{53222400}h^2u^6 + O(u^8), \\
\beta_1 &= \frac{9}{5}h^2 - \frac{19}{1008}h^2u^2 - \frac{83}{151200}h^2u^4 - \frac{1}{98560}h^2u^6 + O(u^8), \\
\beta_2 &= \frac{37}{40}h^2 + \frac{53}{3360}h^2u^2 + \frac{11}{33600}h^2u^4 - \frac{19}{8870400}h^2u^6 + O(u^8), \\
\beta_3 &= \frac{2}{15}h^2 - \frac{11}{5040}h^2u^2 + \frac{17}{151200}h^2u^4 + \frac{173}{13305600}h^2u^6 + O(u^8), \\
\beta_4 &= -\frac{1}{80}h^2 - \frac{31}{20160}h^2u^2 - \frac{67}{604800}h^2u^4 - \frac{109}{17740800}h^2u^6 + O(u^8), \tag{9}
\end{aligned}$$

for y_{n+4} ,

$$\begin{aligned}
 \alpha_0 &= -3, \\
 \alpha_1 &= 4, \\
 \beta_0 &= \frac{2 \sin(wh)h^2w^2 - 7h^2w^2 \sin(2wh) - 6 \sin(wh) - 6 \sin(3wh) + 12 \sin(2wh)}{10 \sin(wh)w^2 + 2w^2 \sin(3wh) - 8w^2 \sin(2wh)}, \\
 \beta_1 &= \frac{35 \sin(wh)h^2w^2 + 7w^2h^2 \sin(3wh) - 4h^2w^2 \sin(2wh) + 4 \sin(wh) + 20 \sin(3wh) - 32 \sin(2wh)}{10 \sin(wh)w^2 + 2w^2 \sin(3wh) - 8w^2 \sin(2wh)}, \\
 \beta_2 &= \frac{w^2h^2 \sin(3wh) - 19h^2w^2 \sin(2wh) - \sin(wh)h^2w^2 - 12 \sin(3wh) + 12 \sin(2wh) + 12 \sin(wh)}{w^2 \sin(3wh) + 5 \sin(wh)w^2 - 4w^2 \sin(2wh)}, \\
 \beta_3 &= \frac{-4h^2w^2 \sin(2wh) - 31 \sin(wh)h^2w^2 - 3w^2h^2 \sin(3wh) + 36 \sin(wh) - 12 \sin(3wh)}{8w^2 \sin(2wh) - 10 \sin(wh)w^2 - 2w^2 \sin(3wh)}, \\
 \beta_4 &= \frac{3h^2w^2 \sin(2wh) + 6 \sin(wh)h^2w^2 + 4 \sin(2wh) - 14 \sin(wh) + 2 \sin(3wh)}{8w^2 \sin(2wh) - 10 \sin(wh)w^2 - 2w^2 \sin(3wh)}.
 \end{aligned} \tag{10}$$

The converted coefficients of the method in series form is given as

$$\begin{aligned}
 \beta_0 &= \frac{9}{40}h^2 + \frac{19}{2016}h^2u^2 + \frac{83}{302400}h^2u^4 + \frac{1}{197120}h^2u^6 + O(u^8), \\
 \beta_1 &= \frac{83}{30}h^2 - \frac{37}{1260}h^2u^2 - \frac{29}{37800}h^2u^4 - \frac{29}{3326400}h^2u^6 + O(u^8), \\
 \beta_2 &= \frac{37}{20}h^2 + \frac{53}{1680}h^2u^2 + \frac{11}{16800}h^2u^4 - \frac{19}{4435200}h^2u^6 + O(u^8), \\
 \beta_3 &= \frac{11}{10}h^2 - \frac{4}{315}h^2u^2 - \frac{1}{9450}h^2u^4 + \frac{1}{69300}h^2u^6 + O(u^8), \\
 \beta_4 &= \frac{7}{120}h^2 + \frac{11}{10080}h^2u^2 - \frac{17}{302400}h^2u^4 - \frac{173}{26611200}h^2u^6 + O(u^8).
 \end{aligned} \tag{11}$$

Equation (2) was differentiated and evaluated at all points which gives the following coefficients and its series form. Differentiating and evaluating at $x = x_n$ gives y'_n .

$$\begin{aligned}
 \widehat{\alpha}_0 &= -\frac{1}{h}, \\
 \widehat{\alpha}_1 &= \frac{1}{h}, \\
 \widehat{\beta}_0 &= \frac{-9h^2w^2 \sin(2wh) + 14 \sin(wh)h^2w^2 - 8wh \cos(3wh)}{32hw^2 \sin(2wh) - 8hw^2 \sin(3wh) - 40hw^2 \sin(wh)} \\
 &\quad + \frac{8wh \cos(2wh) - 16 \sin(2wh) + 8 \sin(3wh) + 8 \sin(wh)}{32hw^2 \sin(2wh) - 8hw^2 \sin(3wh) - 40hw^2 \sin(wh)}, \\
 \widehat{\beta}_1 &= \frac{-20h^2w^2 \sin(2wh) + 27w^2h^2 \sin(3wh) + 7 \sin(wh)h^2w^2}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)} \\
 &\quad + \frac{72wh \cos(3wh) - 24 \cos(wh)hw - 48wh \cos(2wh) + 120 \sin(2wh) - 72 \sin(3wh) - 24 \sin(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)} \\
 &\quad + \frac{72wh \cos(3wh) - 24 \cos(wh)hw - 48wh \cos(2wh) + 120 \sin(2wh) - 72 \sin(3wh) - 24 \sin(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)}, \\
 \widehat{\beta}_2 &= \frac{-11w^2h^2 \sin(3wh) + 11 \sin(wh)h^2w^2 - 7h^2w^2 \sin(2wh) - 36wh \cos(3wh)}{48hw^2 \sin(2wh) - 12hw^2 \sin(3wh) - 60hw^2 \sin(wh)} \\
 &\quad + \frac{36 \cos(wh)hw + 36 \sin(3wh) - 36 \sin(wh) - 36 \sin(2wh)}{48hw^2 \sin(2wh) - 12hw^2 \sin(3wh) - 60hw^2 \sin(wh)}, \\
 \widehat{\beta}_3 &= \frac{-7w^2h^2 \sin(3wh) - 20h^2w^2 \sin(2wh) + 13 \sin(wh)h^2w^2 - 48wh \cos(2wh)}{24hw^2 \sin(3wh) + 120hw^2 \sin(wh) - 96hw^2 \sin(2wh)} \\
 &\quad + \frac{72 \cos(wh)hw - 24wh \cos(3wh) + 24 \sin(3wh) + 24 \sin(2wh) - 120 \sin(wh)}{24hw^2 \sin(3wh) + 120hw^2 \sin(wh) - 96hw^2 \sin(2wh)}, \\
 \widehat{\beta}_4 &= \frac{7h^2w^2 \sin(2wh) - 2 \sin(wh)h^2w^2 + 24wh \cos(2wh) - 24 \cos(wh)hw - 24 \sin(2wh) + 48 \sin(wh)}{24hw^2 \sin(3wh) + 120hw^2 \sin(wh) - 96hw^2 \sin(2wh)}.
 \end{aligned} \tag{12}$$

The converted coefficients of the method in series form is given as

$$\begin{aligned}
 \widehat{\beta}_0 &= -\frac{367}{1440}h - \frac{199}{24192}hu^2 - \frac{1543}{3628800}hu^4 - \frac{281}{9123840}hu^6 + O(u^8), \\
 \widehat{\beta}_1 &= -\frac{3}{8}h + \frac{337}{15120}hu^2 + \frac{457}{453600}hu^4 - \frac{289}{4435200}hu^6 + O(u^8), \\
 \widehat{\beta}_2 &= \frac{47}{240}h - \frac{353}{20160}hu^2 - \frac{19}{40320}hu^4 - \frac{569}{53222400}hu^6 + O(u^8), \\
 \widehat{\beta}_3 &= -\frac{29}{360}h + \frac{1}{945}hu^2 - \frac{43}{113400}hu^4 - \frac{127}{2494800}hu^6 + O(u^8), \\
 \widehat{\beta}_4 &= \frac{7}{480}h + \frac{289}{120960}hu^2 + \frac{139}{518400}hu^4 + \frac{2899}{106444800}h^2u^6 + O(u^8).
 \end{aligned} \tag{13}$$

Differentiating and evaluating at $x = x_n$ gives y'_{n+1} .

$$\begin{aligned}
 \widehat{\alpha}_0 &= -\frac{1}{h}, \\
 \widehat{\alpha}_1 &= \frac{1}{h}, \\
 \widehat{\beta}_0 &= \frac{26 \sin(wh)h^2w^2 - 19h^2w^2 \sin(2wh) - 24 \cos(wh)hw}{120hw^2 \sin(wh) - 96hw^2 \sin(2wh) + 24hw^2 \sin(3wh)} \\
 &\quad + \frac{24wh \cos(2wh) - 24 \sin(wh) + 48 \sin(2wh) - 24 \sin(3wh)}{120hw^2 \sin(wh) - 96hw^2 \sin(2wh) + 24hw^2 \sin(3wh)}, \\
 \widehat{\beta}_1 &= \frac{23 \sin(wh)h^2w^2 - 16h^2w^2 \sin(2wh) + 19w^2h^2 \sin(3wh) + 48 \cos(wh)hw - 72wh \cos(2wh)}{120hw^2 \sin(wh) - 96hw^2 \sin(2wh) + 24hw^2 \sin(3wh)} \\
 &\quad + \frac{24wh + 24 \sin(wh) - 120 \sin(2wh) + 72 \sin(3wh)}{120hw^2 \sin(wh) - 96hw^2 \sin(2wh) + 24hw^2 \sin(3wh)}, \\
 \widehat{\beta}_2 &= \frac{5 \sin(wh)h^2w^2 - 5w^2h^2 \sin(3wh) - 13h^2w^2 \sin(2wh)}{60hw^2 \sin(wh) - 48hw^2 \sin(2wh) + 12hw^2 \sin(3wh)} \\
 &\quad + \frac{36wh \cos(2wh) - 36wh - 36 \sin(3wh) + 36 \sin(wh) + 36 \sin(2wh)}{60hw^2 \sin(wh) - 48hw^2 \sin(2wh) + 12hw^2 \sin(3wh)}, \\
 \widehat{\beta}_3 &= \frac{-16h^2w^2 \sin(2wh) - 3w^2h^2 \sin(3wh) - 7 \sin(wh)h^2w^2}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)} \\
 &\quad + \frac{24wh \cos(2wh) + 48 \cos(wh)hw - 72wh - 24 \sin(2wh) - 24 \sin(3wh) + 120 \sin(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)}, \\
 \widehat{\beta}_4 &= \frac{h^2w^2 \sin(2wh) + 2 \sin(wh)h^2w^2 - 8 \cos(wh)hw + 8wh + 8 \sin(2wh) - 16 \sin(wh)}{32hw^2 \sin(2wh) - 8hw^2 \sin(3wh) - 40hw^2 \sin(wh)}.
 \end{aligned} \tag{14}$$

The converted coefficients of the method in series form is given as

$$\begin{aligned}
 \widehat{\beta}_0 &= \frac{3}{32}h + \frac{731}{120960}hu^2 + \frac{1439}{3628800}hu^4 - \frac{157}{5068800}hu^6 + O(u^8), \\
 \widehat{\beta}_1 &= \frac{47}{90}h - \frac{97}{6048}hu^2 - \frac{853}{907200}hu^4 - \frac{5317}{79833600}hu^6 + O(u^8), \\
 \widehat{\beta}_2 &= -\frac{41}{240}h + \frac{(239)}{20160}hu^2 + \frac{89}{201600}hu^4 + \frac{743}{53222400}hu^6 + O(u^8), \\
 \widehat{\beta}_3 &= \frac{1}{15}h + \frac{1}{4320}hu^2 + \frac{319}{907200}hu^4 + \frac{1277}{26611200}hu^6 + O(u^8), \\
 \widehat{\beta}_4 &= \frac{17}{1440}h - \frac{253}{120960}hu^2 - \frac{181}{725760}hu^4 - \frac{1681}{63866880}h^2u^6 + O(u^8).
 \end{aligned} \tag{15}$$

Differentiating and evaluating at $x = x_n$ gives y'_{n+2} .

$$\begin{aligned}
 \widehat{\alpha}_0 &= -\frac{1}{h}, \\
 \widehat{\alpha}_1 &= \frac{1}{h}, \\
 \widehat{\beta}_0 &= \frac{29h^2w^2 \sin(2wh) - 22 \sin(wh)h^2w^2 - 24wh \cos(wh) + 24wh - 48 \sin(2wh) + 24 \sin(3wh) + 24 \sin(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)}, \\
 \widehat{\beta}_1 &= \frac{29w^2h^2 \sin(3wh) + 113 \sin(wh)h^2w^2 - 28h^2w^2 \sin(2wh) - 48wh \cos(wh) + 48wh + 72 \sin(3wh) + 24 \sin(wh) - 120 \sin(2wh)}{24hw^2 \sin(3wh) + 120hw^2 \sin(wh) - 96hw^2 \sin(2wh)}, \\
 \widehat{\beta}_2 &= \frac{w^2h^2 \sin(3wh) - \sin(wh)h^2w^2 - 19h^2w^2 \sin(2wh) - 12 \sin(3wh) + 12 \sin(wh) + 12 \sin(2wh)}{4hw^2 \sin(3wh) + 20hw^2 \sin(wh) - 16hw^2 \sin(2wh)}, \\
 \widehat{\beta}_3 &= \frac{w^2h^2 \sin(3wh) + 85 \sin(wh)h^2w^2 + 28h^2w^2 \sin(2wh) + 48wh \cos(wh) - 48wh + 24 \sin(3wh) - 120 \sin(wh) + 24 \sin(2wh)}{24hw^2 \sin(3wh) + 120hw^2 \sin(wh) - 96hw^2 \sin(2wh)}, \\
 \widehat{\beta}_4 &= \frac{-34 \sin(wh)h^2w^2 - h^2w^2 \sin(2wh) - 24wh \cos(wh) + 24wh + 48 \sin(wh) - 24 \sin(2wh)}{24hw^2 \sin(3wh) + 120hw^2 \sin(wh) - 96hw^2 \sin(2wh)}. \tag{16}
 \end{aligned}$$

The converted coefficients of the method in series form is given as

$$\begin{aligned}
 \widehat{\beta}_0 &= \frac{97}{1440}h + \frac{1}{640}hu^2 - \frac{439}{3628800}hu^4 - \frac{7211}{319334400}hu^6 + O(u^8), \\
 \widehat{\beta}_1 &= \frac{361}{360}h - \frac{29}{5040}hu^2 + \frac{17}{90720}hu^4 + \frac{1817}{39916800}hu^6 + O(u^8), \\
 \widehat{\beta}_2 &= \frac{37}{80}h + \frac{53}{6720}hu^2 + \frac{11}{67200}hu^4 - \frac{19}{17740800}hu^6 + O(u^8), \\
 \widehat{\beta}_3 &= -\frac{13}{360}h - \frac{1}{210}hu^2 - \frac{23}{56700}hu^4 - \frac{1}{22680}hu^6 + O(u^8), \\
 \widehat{\beta}_4 &= \frac{1}{288}h + \frac{43}{40320}hu^2 + \frac{91}{518400}hu^4 + \frac{7097}{518400}h^2u^6 + O(u^8). \tag{17}
 \end{aligned}$$

Differentiating and evaluating at $x = x_n$ gives y'_{n+3} .

$$\begin{aligned}
 \widehat{\alpha}_0 &= -\frac{1}{h}, \\
 \widehat{\alpha}_1 &= \frac{1}{h}, \\
 \widehat{\beta}_0 &= \frac{9h^2w^2 \sin(2wh) + 2 \sin(wh)h^2w^2 + 8wh \cos(wh) - 8wh - 16 \sin(2wh) + 8 \sin(3wh) + 8 \sin(wh)}{32hw^2 \sin(2wh) - 8hw^2 \sin(3wh) - 40hw^2 \sin(wh)}, \\
 \widehat{\beta}_1 &= \frac{16h^2w^2 \sin(2wh) - 27w^2h^2 \sin(3wh) - 191 \sin(wh)h^2w^2 - 24wh \cos(2wh) - 48wh \cos(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)} \\
 &\quad + \frac{-24 \sin(wh) + 72wh + 120 \sin(2wh) - 72 \sin(3wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)}, \\
 \widehat{\beta}_2 &= \frac{101h^2w^2 \sin(2wh) - 11w^2h^2 \sin(3wh) + 11 \sin(wh)h^2w^2}{48hw^2 \sin(2wh) - 12hw^2 \sin(3wh) - 60hw^2 \sin(wh)} \\
 &\quad + \frac{36wh \cos(2wh) - 36wh - 36 \sin(2wh) + 36 \sin(3wh) - 36 \sin(wh)}{48hw^2 \sin(2wh) - 12hw^2 \sin(3wh) - 60hw^2 \sin(wh)}, \\
 \widehat{\beta}_3 &= \frac{-16h^2w^2 \sin(2wh) - 11w^2h^2 \sin(3wh) - 175 \sin(wh)h^2w^2 - 72wh \cos(2wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)} \\
 &\quad + \frac{48wh \cos(wh) + 24wh - 24 \sin(2wh) - 24 \sin(3wh) + 120 \sin(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)}, \\
 \widehat{\beta}_4 &= \frac{11h^2w^2 \sin(2wh) + 38 \sin(wh)h^2w^2 + 24wh \cos(2wh) - 24wh \cos(wh) + 24 \sin(2wh) - 48 \sin(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)}. \tag{18}
 \end{aligned}$$

The converted coefficients of the method in series form is given as

$$\begin{aligned}\widehat{\beta}_0 &= \frac{119}{1440}h + \frac{571}{120960}hu^2 + \frac{1103}{3628800}hu^4 + \frac{8291}{319334400}hu^6 + O(u^8), \\ \widehat{\beta}_1 &= \frac{9}{10}h - \frac{65}{6048}hu^2 - \frac{517}{907200}hu^4 - \frac{59}{1267200}hu^6 + O(u^8), \\ \widehat{\beta}_2 &= \frac{263}{240}h + \frac{79}{20160}hu^2 - \frac{23}{201600}hu^4 - \frac{857}{53222400}hu^6 + O(u^8), \\ \widehat{\beta}_3 &= \frac{4}{9}h + \frac{167}{30240}hu^2 + \frac{131}{181440}hu^4 + \frac{5431}{79833600}hu^6 + O(u^8), \\ \widehat{\beta}_4 &= -\frac{11}{480}h - \frac{59}{17280}hu^2 - \frac{1241}{3628800}hu^4 - \frac{667}{21288960}h^2u^6 + O(u^8).\end{aligned}\quad (19)$$

Differentiating and evaluating at $x = x_n$ gives y'_{n+4} .

$$\begin{aligned}\widehat{\alpha}_0 &= -\frac{1}{h}, \\ \widehat{\alpha}_1 &= \frac{1}{h}, \\ \widehat{\beta}_0 &= \frac{37h^2w^2 \sin(2wh) + 10 \sin(wh)h^2w^2 + 24wh \cos(2wh) - 24wh \cos(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)} \\ &\quad + \frac{-48 \sin(2wh) + 24 \sin(3wh) + 24 \sin(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)}, \\ \widehat{\beta}_1 &= \frac{-20h^2w^2 \sin(2wh) - 37w^2h^2 \sin(3wh) - 185 \sin(wh)h^2w^2 - 48wh \cos(2wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)} \\ &\quad + \frac{72wh \cos(wh) - 24wh \cos(3wh) + 120 \sin(2wh) - 72 \sin(3wh) - 24 \sin(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)}, \\ \widehat{\beta}_2 &= \frac{121h^2w^2 \sin(2wh) + 5w^2h^2 \sin(3wh) - 5 \sin(wh)h^2w^2 - 36wh \cos(wh)}{48hw^2 \sin(2wh) - 12hw^2 \sin(3wh) - 60hw^2 \sin(wh)} \\ &\quad + \frac{36wh \cos(3wh) - 36 \sin(2wh) + 36 \sin(3wh) - 36 \sin(wh)}{48hw^2 \sin(2wh) - 12hw^2 \sin(3wh) - 60hw^2 \sin(wh)}, \\ \widehat{\beta}_3 &= \frac{20h^2w^2 \sin(2wh) - 57w^2h^2 \sin(3wh) - 205 \sin(wh)h^2w^2 + 48wh \cos(2wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)} \\ &\quad + \frac{24wh \cos(wh) - 72wh \cos(3wh) - 24 \sin(2wh) - 24 \sin(3wh) + 120 \sin(wh)}{96hw^2 \sin(2wh) - 24hw^2 \sin(3wh) - 120hw^2 \sin(wh)}, \\ \widehat{\beta}_4 &= \frac{19h^2w^2 \sin(2wh) - 10 \sin(wh)h^2w^2 + 8wh \cos(3wh) - 8wh \cos(2wh) + 8 \sin(2wh) - 16 \sin(wh)}{32hw^2 \sin(2wh) - 8hw^2 \sin(3wh) - 40hw^2 \sin(wh)}.\end{aligned}\quad (20)$$

The converted coefficients of the method in series form is given as

$$\begin{aligned}\widehat{\beta}_0 &= \frac{9}{160}h + \frac{29}{120960}hu^2 - \frac{31}{145152}hu^4 - \frac{89}{3225600}hu^6 + O(u^8), \\ \widehat{\beta}_1 &= \frac{377}{360}h - \frac{5}{432}hu^2 + \frac{73}{453600}hu^4 + \frac{2089}{39916800}hu^6 + O(u^8), \\ \widehat{\beta}_2 &= \frac{35}{48}h + \frac{671}{20160}hu^2 + \frac{23}{28800}hu^4 + \frac{13}{1520640}hu^6 + O(u^8), \\ \widehat{\beta}_3 &= \frac{161}{120}h - \frac{31}{945}hu^2 - \frac{139}{113400}hu^4 - \frac{53}{831600}hu^6 + O(u^8), \\ \widehat{\beta}_4 &= \frac{469}{1440}h + \frac{1313}{120960}hu^2 + \frac{1741}{3628800}hu^4 + \frac{9721}{319334400}h^2u^6 + O(u^8).\end{aligned}\quad (21)$$

3. Basic properties of the method

The implicit schemes from equation (6)-(13) derived are discrete schemes belonging to the class of LMMs of the form

$$\sum_{j=0}^k \alpha_j y_{n+j} = h^2 \sum_{j=0}^k \beta_j f_{n+j}. \quad (22)$$

Following Refs. [2, 11], we define the Local Truncation Error(LTE) associated with equation (22) by difference operator;

$$L[y(x); h] = \sum_{j=0}^k [\alpha_j y(x + jh) - h^2 \beta_j y''(x + jh)]. \quad (23)$$

Expanding (23) by Taylor series, we have

$$L[y(x); h] = C_0 y(x) + C_1 h y'(x) + \cdots + C_q h^q y^{(q)}(x) + \cdots, \quad (24)$$

where

$$\begin{aligned} C_0 &= \alpha_0 + \alpha_1 + \alpha_2 + \cdots + \alpha_k, \\ C_1 &= \alpha_1 + 2\alpha_2 + \cdots + k\alpha_k, \\ C_2 &= \frac{1}{2!}(\alpha_1 + 2^2\alpha_2 + \cdots + k^2\alpha_k) - (\beta_0 + \beta_1 + \beta_2 + \cdots + \beta_k), \\ &\vdots \\ C_p &= \frac{1}{p!}(\alpha_1 + 2^p\alpha_2 + \cdots + k^p\alpha_k) - \frac{1}{(q-2)!} \\ &\quad (\beta_1 + 2^{q-2}\beta_2 + \cdots + k^{q-2}\beta_k), q \geq 3. \end{aligned}$$

3.1. Order and error constant

The LMM (22) is said to be of order p if $C_0 = C_1 = C_2 = \cdots = C_p = C_{p+1} = 0$ and $C_{p+2} \neq 0$ is the error constant. Equation (6) has order $p = 5$ and error constant given by $C_{p+2} = \frac{1}{240}$. Equation (8) has order $p = 5$ and error constant given by $C_{p+2} = \frac{1}{120}$, Equation (10) has order $p = 5$ and error constant given by $C_{p+2} = \frac{1}{120}$ and Equation (12) has order $p = 5$ and error constant given by $C_{p+2} = \frac{-191}{720}$.

3.2. Zero stability

Theorem 1. Zero-stability [1, 11]: A block method is said to be zero stable if as $h \rightarrow 0$, the roots $r_j, j = 1(1)k$ of the first characteristic polynomials $\rho(r) = 0$ that is

$$\rho(r) = \det \left[\sum A^{(0)} R^{k-1} \right] = 0,$$

satisfying $|R| \leq 1$, must be simple.

$$\rho(z) = \det[zA^{(0)} - A^{(i)}] = 0,$$

$$\left[z \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \right] = 0$$

$$\left[\begin{pmatrix} z & 0 & 0 & 0 \\ 0 & z & 0 & 0 \\ 0 & 0 & z & 0 \\ 0 & 0 & 0 & z \end{pmatrix} - \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \right] = 0$$

$$z^3(z-1) = 0, \quad z = 1, 0, 0, 0$$

since $|z = 1, 0, 0, 0| \leq 1$, the method is stable.

3.3. Consistency

According to Ref. [12], a numerical method is said to be consistent if the following conditions are satisfied.

- (i) The order of the method must be greater or equal to 1. i.e. $p \geq 1$;
- (ii) $\sum_{j=0}^k \alpha_j = 0$;
- (iii) $\rho(r) = \rho'(r) = 0$;
- (iv) $\rho''(r) = 2!\sigma(r)$.

where ρ and σ are the first and second characteristics polynomial of equation (5). When these conditions are applied to the main scheme, it is found to be consistent.

Table 1: Numerical results and the absolute error for test problem 1.

x	Exact	Computed	Error
0.1	-0.10517091807564762	-0.10517091807437191	1.27571E-12
0.2	-0.22140275816016983	-0.22140275815480269	5.36714E-12
0.3	-0.34985880757600310	-0.34985880756315065	1.285245E-11
0.4	-0.49182469764127032	-0.49182469761687007	2.440025E-11
0.5	-0.64872127070012815	-0.64872127065934589	4.078226E-11
0.6	-0.82211880039050897	-0.82211880032762140	6.288757E-11
0.7	-1.01375270747047652	-1.01375270737873767	9.173885E-11
0.8	-1.22554092849246760	-1.22554092836395704	1.2851056E-10
0.9	-1.45960311115694966	-1.45960311098240009	1.7454957E-10
1.0	-1.71828182845904524	-1.71828182822764679	2.3139845E-10

Table 2: The comparison of error in new method and some existing methods for test problem 1.

x	Error in new method	Error in Ref. [13]	Error in Ref. [14]	Error in Ref. [15]
0.1	1.27571E-12	-	6.899835E-11	4.462679E-11
0.2	5.36714E-12	3.46017000E-09	1.525099E-10	9.864032E-11
0.3	1.285245E-11	5.67600300E-09	2.528244E-10	1.635218E-10
0.4	2.440025E-11	7.64127000E-09	3.725524E-10	2.409591E-10
0.5	4.078226E-11	1.04971347E-08	5.146680E-10	3.328765E-10
0.6	6.288757E-11	1.44950355E-08	6.825557E-10	4.414623E-10
0.7	9.173885E-11	1.87822380E-08	8.800651E-10	5.692067E-10
0.8	1.2851056E-10	2.27988702E-08	1.111569E-09	7.189380E-10
0.9	1.7454957E-10	2.82582100E-08	1.382034E-09	8.938681E-10
1.0	2.3139845E-10	3.55473540E-08	1.697095E-09	1.097642E-09

Table 3: Numerical results and the absolute error for test problem 2.

x	Exact	Computed	Error
0.01	1.01979867335991086	1.01979867335968196	2.2890E-13
0.02	1.03918944084761210	1.03918944084696135	6.5075E-13
0.03	1.05816454641464877	1.05816454641402239	6.2638E-13
0.04	1.07671640027179207	1.07671640027015420	1.63787E-12
0.05	1.09483758192485392	1.09483758192097119	3.88273E-12
0.06	1.11252084314278561	1.11252084313620862	6.57699E-12
0.07	1.12975911085687365	1.12975911084716780	9.70585E-12
0.08	1.14654548998987291	1.14654548997661875	1.325416E-12
0.09	1.16287326621394559	1.16287326619673912	1.720647E-11
0.1	1.17873590863630285	1.17873590861475587	2.154698E-11

3.4. Convergence

Theorem 2. [11]: The necessary and sufficient condition for a linear multistep method to be convergent is for it to be consistent and zero stable. Thus our block method is convergent since it is zero stable and consistent.

4. Implementation and numerical examples

This section examines the performance of the new method by applying it to some second-order ODEs . The test problems outcomes are displayed in tabular form.

Error= |exact solution – computed solution|.

Test Problem 1:

$$y'' = y', \quad y(0) = 1, \quad y'(0) = 0, \quad h = 0.1.$$

Exact solution: $y(x) = 1 - \exp(x)$.

Table 4: The comparison of error in new method and some existing methods for test problem 2.

x	Error in new method	Error in Ref. [16]
0.01	2.2890E-13	3.409E-11
0.02	6.5075E-13	3.239E-11
0.03	6.2638E-13	3.465E-11
0.04	1.63787E-12	2.40E-13
0.05	3.88273E-12	1.780E-12
0.06	6.57699E-12	7.467E-11
0.07	9.70585E-12	3.904E-11
0.08	1.325416E-12	4.132E-11
0.09	1.720647E-11	1.197E-10
0.1	2.154698E-11	8.342E-11

Table 5: Numerical results and the absolute error for test problem 3.

x	Exact	Computed	Error
0.01	0.904837418035959573	0.904837419827875764	1.791916191E-9
0.02	0.818730753077981859	0.818730757391773436	4.313791577E-9
0.03	0.740818220681717866	0.740818227549595599	6.867877733E-9
0.04	0.670320046035639301	0.670320054867431370	8.831792069E-9
0.05	0.606530659712633424	0.606530682641211184	2.2928577760E-8
0.06	0.548811636094026433	0.548811675353608366	3.9259581933E-8
0.07	0.496585303791409515	0.496585361588883748	5.7797474233E-8
0.08	0.449328964117221591	0.449329042672387956	7.8555166365E-8
0.09	0.406569659740599112	0.406569761324812764	1.01584213652E-7
0.1	0.367879441171442322	0.367879568145210102	1.26973767780E-7

Table 6: The comparison of error in new method and some existing methods for test problem 3.

x	Error in new method	Error in Ref. [17]
0.01	1.791916191E-9	1.29E-8
0.02	4.313791577E-9	3.01E-8
0.03	6.867877733E-9	5.04E-8
0.04	8.831792069E-9	9.32E-10
0.05	2.2928577760E-8	1.40E-7
0.06	3.9259581933E-8	1.90E-7
0.07	5.7797474233E-8	2.58E-7
0.08	7.8555166365E-8	3.32E-7
0.09	1.01584213652E-7	-
0.1	1.26973767780E-7	-

Test Problem 2:

$$y'' + \lambda^2 y = 0, \quad y(0) = 1, \quad y'(0) = 2, \quad \lambda = 2.$$

Exact solution: $y(x) = \cos(2x) + \sin(2x)$.

Test Problem 3:

$$y'' - 100y = 0, \quad y(0) = 1, \quad y'(0) = -10, \quad h = 0.01.$$

Exact solution: $y(x) = \exp(-10x)$.

Table 1 shows the results generated for test problem 1 using the proposed method. Table 2 shows the comparison of the results of our method with that of Refs. [13–15]. It is obvious that our method is better in accuracy and also efficient than Refs. [13, 13, 15].

Table 3 represents the outcomes produced for test problem 2 utilizing the proposed method. Meanwhile, Table 4 demonstrates a favourable comparison of the proposed method with the one in Ref. [16].

Table 5 presents the results obtained for test problem 3 using the proposed method. Table 6 shows the comparison of the proposed method with Ref. [17]. It is observed that our method is more efficient and accurate than that of Ref. [17].

The implementation of the new scheme on three specific examples demonstrates its favourable comparison with exact solutions in Table 2 and Table 6 while the result in Table 4 competes well with the existing results.

5. Conclusion

Utilizing collocation and interpolation techniques, a novel class of continuous second derivative block methods for solving ODEs is constructed in this work. This innovative approach combines polynomial and trigonometric functions, implemented through code written in Maple, to develop an approximation solution. The resulting block techniques exhibit continuous coefficients and possess key properties of consistency, zero stability, and convergence. These characteristics contribute to a robust methodology that ensures reliability in solving ODEs, presenting a promising avenue for further exploration and application in this field.

References

- [1] J. D. Lambert, *Computational methods in ODEs*, John Wiley, New York, 1973. <https://doi.org/10.1002/zamm.19740540726>.
- [2] J. D. Lambert, *Numerical methods for ordinary differential systems*. John Wiley, New York, 1991. <https://www.wiley.com/en-be/9780471929901>.
- [3] D. O. Awoyemi, "A new sixth order algorithms for general second order ordinary differential equation", *Inter. J. Computer Math.* **77** (2001) 117. <https://dx.doi.org/10.1080/00207160108805054>.
- [4] S. J. Kayode, "An efficient zero-stable numerical method for fourth-order differential equations", *International Journal of Mathematics and Mathematical Sciences* (2008) 364021. <https://doi.org/10.1155/2008/364021>.
- [5] O. O. Olanegan, B. G. Ogunware & C. O. Alakofa, "Implicit hybrid points approach for solving general second order ordinary differential equations with initial values", *Journal of Advances in Mathematics and Computer Science* **27** (2018) 1. <https://doi.org/10.9734/JAMCS/2018/40447>.
- [6] Z. Omar & J. O. Kuboye, "Application of order nine block method for solving second order ordinary differential equations directly", *Research Journal of Applied Sciences, Engineering and Technology* **1** (2015) 19. <https://doi.org/10.19026/rjaset.11.1671>.
- [7] P. Onumanyi, U. W. Serisena & S. N. Jator, "Continuous finite difference approximation for solving differential equations", *International Journal of Computer Mathematics* **72** (1999) 15. <https://doi.org/10.1080/00207169908804831>.
- [8] A. M. Badmus, "A new eight order implicit block algorithms for the direct solution of second order ordinary differential equations", *American Journal of Computational Mathematics* **4** (2014) 376. <https://doi.org/10.4236/ajcm.2014.44032>.
- [9] S. N. Jator & J. Li, "A self-stationary linear multistep method for a direct solution of general second order IVPs", *International Journal of Computer Mathematics* **86** (2009) 827. <https://doi.org/10.1080/00207160701708250>.
- [10] B. G. Ogunware, F. M. Okafor, E. O. Omole & F. C. Awoyemi, "Solution of second order ordinary differential equations with a one-step hybrid numerical model", *KIU Journal of Science, Engineering and Technology* **02** (2023) 45. <https://doi.org/10.59568/KJSET-2023-2-1-07>.
- [11] S. O. Fatunla, *Numerical methods for initial value problems in ordinary differential equations*, Academic press inc. Harcourt Brace Jovanovich Publishers, New York, 1988. <https://doi.org/10.1016/c22013-0-10643-5>.
- [12] F. F. Ngwane & S. N. Jator, "A family of trigonometrically fitted enright second derivative methods for stiff and oscillatory initial value problems", *Journal of Applied Mathematics* (2015) 343295. <https://doi.org/10.1155/2015/343295>.
- [13] S. J. Kayode & J. O. Adegboro, "Predictor-corrector linear multistep method for direct solution of initial value problems of second order ordinary differential equations", *Asian Journal of Physical and Chemical Sciences* **6** (2018) 1. <https://doi.org/10.9734/AJOPACS/2018/41034>.
- [14] J. O. Kuboye, B. G. Ogunware, O. E. Abolarin & C. O. Mmaduakor, "Single numerical algorithm developed to solving first and second orders ordinary differential equations", *International Journal of Mathematics in Operational Research* **21** (2022) 466. <https://doi.org/10.1504/IJMOR.2021.10038999>.
- [15] B. G. Ogunware, J. O. Kuboye, O. E. Abolarin & C. O. Mmaduakor, "Half-step numerical model of order ten for solution of first and second orders ordinary differential equations", *Journal of Interdisciplinary Mathematics* **25**(2022) 1023. <https://doi.org/10.1080/09720502.2021.1887620>.
- [16] E. O. Omole & B. G. Ogunware, "3- Point single hybrid block method (3PSHBM) for direct solution of general second order initial value problem of ordinary differential equations", *Asian Journal of Scientific Research and Reports* **20** (2018) 1. <https://doi.org/10.9734/JSRR/2018/19862>.
- [17] F. Obarhua & J. O. Adegboro, "An order four continuous numerical method for solving general second order ordinary differential equations", *Journal of the Nigerian Society of Physical Sciences* **3** (2021) 150. <https://doi.org/10.46481/jnsps.2021.150>.