



A Numerical Method using Collocation Approach for the Solution of Volterra-Fredholm Integro-Differential Equations

G. Ajileye*, F. A. Aminu

Department of Mathematics and Statistics, Federal University Wukari, Taraba State, Nigeria

Abstract

This paper consider collocation approach for the numerical solution of Volterra-Fredholm Integro-differential equation using collocation method. We transformed the problem into a system of linear algebraic equations and matrix inversion is adopted to solve the algebraic equations. We substituted the solution algebraic equations into the approximate equation to obtain the numerical result. Some numerical problems are solved to demonstrate the efficiency and consistency of the method.

DOI:10.46481/asr.2022.1.3.58

Keywords: Collocation, Volterra-Fredholm, Integro-differential, Power series, Approximate solution

Article History :

Received: 05 September 2022

Received in revised form: 23 November 2022

Accepted for publication: 28 November 2022

Published: 29 December 2022

© 2022 The Author(s). Published by the Nigerian Society of Physical Sciences under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Communicated by: Tolulope Latunde

1. Introduction

Recently, research has focused on the most effective way to solve integral and integro-differential equations where the unknown function appears under the integral sign. Integro-differential equations are a common component of mathematical descriptions of physical phenomena. These equations can be found in a wide variety of contexts including fluid dynamics, biological models, and chemical kinetics. [1]

Many different approaches have been adopted to investigate the solution of integro-differential equations such as Adomian decompositions method by [2–4], Collocation method by [5–8], Hybrid linear multistep method [9],

*Corresponding author tel. no: +234 8034906427

Email address: ajileye@fuwukari.edu.ng (G. Ajileye)

Chebyshev-Galerkin method [10], Bernoulli matrix method [11], Differential transform method [12], Pseudospectral method [13], Bernstein polynomials method [14]. The Mellin transform approach [15], Homotopy Perturbation [16] and Perturbed method [17]. [18] developed a numerical method for solving the second kind of linear Volterra-Fredholm integro-differential equations using Bernstein polynomial. By using this method, the integro-differential equations are converted into an algebraic equation system and some numerical examples are offered. [19] proposed solution of Fractional Integro-differential Equations of Fredholm type using Legendre Galerkin method. The concept of Legendre Galerkin Method was implemented on some examples of fractional integro-differential equations of Fredholm type to illustrate the practicability of the method.

A numerical methodology based on quartic weighted polynomials for finding the solution of Fractional Integro-Differential Equations (FIDEs) was presented by [20]. The suggested method involved the application of the homotopy perturbation method and the initial approximation as the constructed orthogonal polynomials. [21] presented Akbari-Ganji's Method (AGM) to solve Volterra Integro-Differential Difference Equations (VIDDE) using Legendre polynomials as basis functions. A trial solution function of unknown constants that conform with the differential equations together with the initial conditions were assumed and substituted into the equations under consideration. Collocation approach for the computational solution of Fredholm-Volterra Fractional order of integro-differential equations was presented by [22]. They obtained the linear integral form of the problem and transformed it into a system of linear algebraic equations using standard collocation points. [23] considered first order Volterra Integro-differential equations using standard collocation method. An assumed approximate solution in terms of the constructed polynomial was substituted into the class of integro-differential equation considered. The resulted equation was collocated at appropriate points within the interval of consideration $[0,1]$ to obtain a system of algebraic linear equations.

In this paper, we consider Volterra-Fredholm Integro-differential equation of the form:

$$y'(x) = f(x) + \int_0^x k_1(x,t)y(t)dt + \int_0^1 k_2(x,t)y(t)dt \quad (1)$$

subject to initial condition

$$y(a) = 0 \quad (2)$$

$a \leq x \leq b$, $y(x)$ is the unknown function, $k_1(x,t)$ and $k_2(x,t)$ are the Volterra and Fredholm integral kernel function respectively. $f(x)$ is the known function

2. Basic Definitions

In this section, we provide certain definitions and fundamental ideas for the formulation of the specified problem.

Definition 1: Let (a_n) , $n \geq 0$ be a sequence of real numbers. The power series in x with coefficients a_n is an expression.

$$y(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + \cdots + a_Nx^N = \sum_{n=0}^N a_n x^n = \phi(x) \mathbf{A} \quad (3)$$

where $\phi(x) = [1 \ x \ x^2 \ \cdots \ x^N]$, $\mathbf{A} = [a_0 \ a_1 \ \cdots \ a_N]^T$
then $y(x,n) = x^n \mathbf{A}$, $n = 0(1)N$, $n \in \mathbb{Z}^+$

Definition 2: The desired collocation points within an interval are determined using this method. i.e. $[a,b]$ and is provided by

$$x_i = a + \frac{(b-a)i}{N}, i = 1, 2, 3, \dots, N \quad (4)$$

3. Methodology

In this section, combination of collocation method and power series approximation are employed for the numerical solution of volterra-fredholm integro-differential equations.

Let the solution to (1) and (3) be approximated by

$$y(x) = \phi(x) \mathbf{A} \quad (5)$$

$\phi(x)$ is an interpolating polynomial and $\mathbf{A}$ are parameters to be determined,

$$\phi(x) = \begin{bmatrix} \phi_0(x) & \phi_1(x) & \phi_2(x) & \cdots & \phi_N(x) \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} a_0 & a_1 & a_2 & \cdots & a_N \end{bmatrix}^T$$

substituting (3) into (1) gives

$$\phi'(x)\mathbf{A} = f(x) + \int_0^x k_1(x,t)\phi(t)\mathbf{A}dt + \int_0^1 k_2(x,t)\phi(t)\mathbf{A}dt \quad (6)$$

collecting the like terms

$$\left(\phi'(x) - \int_0^x k_1(x,t)\phi(t)dt + \int_0^1 k_2(x,t)\phi(t)dt \right) \mathbf{A} = f(x) \quad (7)$$

Equation (7) can be writing in this form

$$U(x)\mathbf{A} = f(x) \quad (8)$$

where

$$U(x) = \left(\phi'(x) - \int_0^x k_1(x,t)\phi(t)dt + \int_0^1 k_2(x,t)\phi(t)dt \right)_{1 \times [N+1]}$$

Collocating (8) using the standard collocation points

$$x_i = a + \frac{b-a}{N}i$$

$$U(x_i)\mathbf{A} = f(x_i) \quad (9)$$

where

$$U(x_i) = \begin{bmatrix} U_0(x_0) & U_1(x_0) & U_2(x_0) & \cdots & U_N(x_0) \\ U_0(x_1) & U_1(x_1) & U_2(x_1) & \cdots & U_N(x_1) \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ U_0(x_N) & U_1(x_N) & U_2(x_N) & \cdots & U_N(x_N) \end{bmatrix}, f(x_i) = \begin{bmatrix} f(x_0) \\ f(x_1) \\ \vdots \\ f(x_N) \end{bmatrix}$$

Using the initial conditions

$$y(a) = 0 \quad (10)$$

hence, (10) becomes

$$\phi(a) = 0 \quad (11)$$

Substituting (11) into equation (9) gives

$$U^*(x_i)\mathbf{A} = f^*(x_i) \quad (12)$$

The values of the unknown are solved using matrix inversion. Substituting the values of a_i , $i = 0, 1, 2, \dots, N$ obtained into the approximate solution gives the numerical solution

$$y(x) = \phi(x)U^{*-1}(x_i)f^*(x_i) \quad (13)$$

4. Numerical Examples

In this section, two numerical examples with initial conditions are presented to confirm the efficiency and accuracy of the method. Let $y_n(x)$ and $y(x)$ be the approximate and exact solution respectively. $\text{Error}_N = |y_n(x) - y(x)|$

Example 1: [18] Considering volterra-fredholm integro-differential equation

$$y'(x) = f(x) - \int_0^x x^2 t y(t) dt - \int_0^1 x(x-t) y(t) dt$$

subject to initial conditions

$$y(0) = 1, \quad 0 \leq x \leq 1$$

where

$$f(x) = -\sin(x) + x^2 \cos(x) + x^3 \sin(x) - x^2 + x^2 \sin(1) - x \cos(1) - x \sin(1) + x$$

Exact solution $y(x) = \cos(x)$

Solution 1

We solve this problem at $N = 5, 6$ and 10 but we use $N = 4$ for demonstration

Using approximate solution (3) on example 1 gives

$$\begin{aligned} \phi'(x) \mathbf{A} &= f(x) - \int_0^x x^2 t \phi(t) \mathbf{A} dt - \int_0^1 x(x-t) \phi(t) \mathbf{A} dt \\ \left[\phi'(x) + \int_0^x x^2 t \phi(t) dt + \int_0^1 x(x-t) \phi(t) dt \right] \mathbf{A} &= f(x) \end{aligned} \quad (14)$$

Equation (14) gives

$$U(x) \mathbf{A} = f(x) \quad (15)$$

where

$$U(x) = \phi'(x) + \int_0^x x^2 t \phi(t) dt + \int_0^1 x(x-t) \phi(t) dt$$

collocating at $x_5 = \left[0 \quad \frac{1}{5} \quad \frac{2}{5} \quad \frac{3}{5} \quad \frac{4}{5} \quad 1 \right]$ and substituting the initial conditions gives

$$U(x_i)^* \mathbf{A} = f(x_i)^* \quad (16)$$

where

$$U(x_i) = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ -\frac{37}{625} & \frac{5959}{6250} & \frac{17032}{46875} & \frac{140629}{1562500} & \frac{7813}{1171875} & -\frac{1140619}{82031250} \\ -\frac{17}{2969} & \frac{2969}{70721} & \frac{70721}{172003} & \frac{172003}{86501} & \frac{86501}{571648} & \frac{571648}{17054991} \\ -\frac{625}{78} & \frac{3125}{6287} & \frac{93750}{36927} & \frac{390625}{1649373} & \frac{390625}{327656} & \frac{5859375}{17054991} \\ \frac{625}{278} & \frac{6250}{3633} & \frac{31250}{78697} & \frac{1562500}{766384} & \frac{390625}{2426518} & \frac{27343750}{84473932} \\ \frac{625}{1} & \frac{3125}{2} & \frac{46875}{7} & \frac{390625}{13} & \frac{1171875}{21} & \frac{41015625}{31} \\ 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$f(x_i) = \left[0 \quad -0.2405731317 \quad -0.3951997680 \quad -0.4316932976 \quad -0.3110546806 \quad 0 \quad 1 \right]^T$$

We now solve for the unknown values $\mathbf{A}$ (16) making use of matrix inversion results in;

$$y_5 = \begin{pmatrix} 1.000000014500000 - 0.1100900000e - 5x - 0.499665406900000x^2 \\ -0.1934020400000e - 2x^3 + 0.45850805300000e - 1x^4 - 0.3948219000000e - 2x^5 \end{pmatrix}$$

Applying the same procedure for $N = 6$ and 10 gives

$$y_6 = \begin{pmatrix} 0.999999999900000 - 1.094862422900000 \times 10^{-8}x - 0.499989658700000x^2 \\ -0.79348700000e - 4x^3 + 0.41903091000000e \\ -1x^4 - 0.323985800000e - 3x^5 - 0.1207684000000e - 2x^6 \end{pmatrix}$$

Table 1. Exact and approximate values, Example 1

x	Exact	$N = 5$	$N = 6$	$N = 10$
0.2	0.980066577841242	0.980069803739200	0.980066640551043	0.980066578523643
0.4	0.921060994002885	0.921062682583520	0.921061026653094	0.921060999303860
0.6	0.825335614909678	0.825337309927040	0.825335678632713	0.825335614259620
0.8	0.696706709347165	0.696709792368160	0.696706738330861	0.696706709330861
1.0	0.540302305868140	0.540302072600000	0.540302402751376	0.540302305851376

Table 2. Absolute Error for Example 1

x	error ₅	error ₆	error ₁₀
0.2	0.000003225897958	6.27098010000000001e-8	7.2364300000000000e-10
0.4	0.000001688580635	3.2650209000000000e-8	5.3009750000000000e-9
0.6	0.000001695017362	6.3723035000000000e-8	1.6465005800000000e-9
0.8	0.000003083020995	2.8983696000000000e-8	1.8604468190000000e-9
1.0	2.3326814000000000e-7	9.6883236000000000e-8	1.0605826775000000e-9

$$y_{10} = \begin{pmatrix} 1.000000000000000 - 5.144329406903130 \times 10^{-12}x - 0.499999944586307x^2 \\ -9.462237358093260 \times 10^{-7}x^3 + 0.41672945022583e - 1x^4 - 0.18835067749e - 4x^5 \\ -0.1343965530396e - 2x^6 - 0.78201293945e - 4x^7 \\ +0.87261199951e - 4x^8 - 0.31232833862e - 4x^9 + 0.4619359970e - 5x^{10} \end{pmatrix}$$

Example 2 [18] Considering fractional integro-differential equation

$$y'(x) = f(x) + \int_0^x y(t)dt + \int_0^1 y(t)dt$$

subject to initial conditions

$$y(0) = 0, \quad 0 \leq x \leq 1$$

where

$$f(x) = 2e^x - 2$$

Exact solution $y(x) = xe^x$

Solution 2

Using approximate solution (3) on example 2 gives

$$\phi'(x) \mathbf{A} = f(x) + \int_0^x \phi(t) \mathbf{A} dt + \int_0^1 \phi(t) \mathbf{A} dt$$

$$\left[\phi'(x) - \int_0^x \phi(t) dt - \int_0^1 \phi(t) dt \right] \mathbf{A} = f(x) \quad (17)$$

Equation (17) gives

$$U(x) \mathbf{A} = f(x) \quad (18)$$

where

$$U(x) = \phi'(x) - \int_0^x \phi(t) dt - \int_0^1 \phi(t) dt$$

collocating at $x_4 = \left[0 \quad \frac{1}{4} \quad \frac{2}{4} \quad \frac{3}{4} \quad 1 \right]$ and substituting the initial conditions gives

$$U(x_i)^* \mathbf{A} = f(x_i)^* \quad (19)$$

Table 3. Exact and approximate values, Example 2

x	Exact	$N = 4$	$N = 6$	$N = 10$
0.2	0.244280551632034	0.245107750566580	0.244280474366985	0.244280551735439
0.4	0.596729879056508	0.597952145575241	0.596729787759204	0.596729879059536
0.6	1.093271280234310	1.094251609446400	1.093271159740840	1.093271280740840
0.8	1.780432742793980	1.781874552875250	1.780432577181920	1.7804327427932846
1.0	2.718281828459050	2.720832262494860	2.718281685733800	2.7182818284583942

Table 4. Absolute Error for Example 2

x	error ₄	error ₆	error ₁₀
0.2	0.000827198934546	7.7265049000000000e-8	1.0103405000000000e-10
0.4	0.001222266518733	9.1297304000000000e-8	1.0903028000000000e-10
0.6	0.000980329212090	1.2049347000000000e-7	2.6958374929600000e-9
0.8	0.001441810081270	1.6561206000000000e-7	1.0116729458000000e-9
1.0	0.002550434035810	1.4272525000000000e-7	3.5968446284000000e-9

where

$$U(x_i) = \begin{bmatrix} -1 & \frac{1}{2} & -\frac{1}{3} & -\frac{1}{4} & -\frac{1}{5} \\ -\frac{5}{4} & \frac{15}{32} & \frac{31}{192} & -\frac{65}{1024} & -\frac{141}{1024} \\ -\frac{3}{2} & \frac{5}{8} & \frac{5}{96} & \frac{31}{512} & \frac{47}{128} \\ -\frac{7}{4} & \frac{7}{32} & \frac{197}{192} & \frac{1391}{1024} & \frac{7373}{5120} \\ -2 & 0 & \frac{4}{3} & \frac{5}{2} & \frac{18}{5} \\ 1 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$f(x_i) = \begin{bmatrix} 0 & 0.568050833375 & 1.29744254140 & 2.23400003322534 & 3.4365636569181 & 0 \end{bmatrix}^T$$

We now solve for the unknown values $\mathbf{A}$ (19) making use of matrix inversion results in;

$$y_4 = \begin{pmatrix} -0.7110336836e - 5 + 1.000614432260340x \\ +1.039187111971780x^2 + 0.364817101049653x^3 + 0.316220727549933x^4 \end{pmatrix}$$

Applying the same procedure for $N = 7$ and 10 gives

$$y_6 = \begin{pmatrix} -1.728131526768100e - 14 + 0.999999894930511x \\ +0.999982466411893x^2 + 0.500181985087693x^3 + 0.165893773548305x^4 \\ +0.43341163545847e - 1x^5 + 0.6405057851225e - 2x^6 + 0.2477344358340e - 2x^7 \end{pmatrix}$$

$$y_{10} = \begin{pmatrix} -7.646335821454020e - 15 + 0.99999999883585x + 0.999999657273293x^2 \\ +0.500003099441528x^3 + 0.166658401489258x^4 + 0.41748046875000e - 1x^5 \\ +0.008087158203125x^6 + 0.001541137695313x^7 + 0.000213623046875x^8 \\ +0.000007629394531x^9 - 0.000005722045898x^{10} \end{pmatrix}$$

5. Conclusion

Volterra-Fredholm integro-differential equations are complex to solve analytically. Hence, approximate solution is needed. This paper considered collocation method for the solution of volterra-fredholm integro-differential equations. The method is consistent, efficient and easy to compute. The results of example 1 as shown in the table 1 and 2 shows the approximate solutions converges to the exact solution when the values of N increases. In example 2, The approximate solution at $N = 4$ gives $y_4(x) = -0.7110336836e - 5 + 1.000614432260340x + 1.039187111971780x^2 + 0.364817101049653x^3 + 0.316220727549933x^4$ and solving for $N=6$ and 10, we obtained Table 3 which display the

results obtained at $x = 0.2$ to 1.0 for various values of N and the exact solution. Error of example 2 as shown in Table 4 indicates that as the values of N increases, the error becomes smaller. All computations are done with the aid of Maple 18.

References

- [1] A. M. Wazwaz & S. M. El-Sayed, “A new modification of the Adomian decomposition method for linear and nonlinear operators”, (Doctoral dissertation). App. Math. Comput. **181** (2001) 393.
- [2] D. A. Gegele, O. P. Evans & D. Akoh, “Numerical solution of higher order linear Fredholm integro-differential equations”, American Journal of Engineering Research **3** (2014) 243.
- [3] R. H. Khan & H. O. Bakodah, “Adomian decomposition method and its modification for nonlinear Abel’s integral equations”, Computers and Mathematics with Applications **7** (2013) 2349.
- [4] R. C. Mittal & R. Nigam, “Solution of fractional integro-differential equations by Adomian decomposition method”, The International Journal of Applied Mathematics and Mechanics **2** (2008) 87.
- [5] A. O. Adesanya, Y. A. Yahaya, B. Ahmed & R. O. Onsachi, “Numerical Solution of Linear integral and Integro-Differential Equations Using Boubakar Collocation Method”, International Journal of Mathematical Analysis and Optimization: Theory and Application **2** (2019) 592.
- [6] A. O. Agbolade & T. A. Anake, “Solution of first order volterra linear integro differential equations by collocation method”, J. Appl. Math., Article ID. 1510267 (2017). <https://doi.org/10.1155/2017/1510267>
- [7] S. Nemati, P. Lima & Y. Ordokhani, “Numerical method for the mixed Volterra-Fredholm integral equations using hybrid Legendre function”, Conference Application of Mathematics (2015) 184.
- [8] G. Mehdiyera, M. Imanova & V. Ibrahim, “Solving Volterra integro differential equation by second derivative methods”, 43rd Appl. Math. Inf. Sci. **9** (2015) 2521.
- [9] G. Mehdiyeva, V. Ibrahimov & M. Imanova, “On the Construction of the Multistep Methods to Solving the Initial-Value Problem for ODE and the Volterra Integro-Differential Equations”, IAPE, Oxford, United Kingdom (2019), ISBN: 978-1-912532-05-6.
- [10] K. Issa & F. Saleh, “Approximate solution of perturbed Volterra Fredholm integro differential equation by Chebyshev-Galerkin method”, Journal of Mathematics (2017). <https://doi.org/10.1155/2017/8213932>.
- [11] A. H. Bhrawy, E. Tohidi & F. Soleymani, “A new Bernoulli matrix method for solving high order linear and nonlinear Fredholm integro-differential equations with piecewise interval”, Appl. Math. Comput. **219** (2012) 482.
- [12] C. Ercan & T. Kharerah, “Solving a class of Volterra integral system by the differential transform method”, Int. J. Nonlinear Sci. **16** (2013) 87.
- [13] M. El-kady & M. Biomy, “Efficient Legendre pseudospectral method for solving integral and integro differential equation”, Commom Non-linear Sci. Numer Simulat (2010) 1724.
- [14] N. Irfan, S. Kumar & S. Kapoor, “Bernstein Operational Matrix Approach for Integro-Differential Equation Arising in Control theory”, Nonlinear Engineering **3** (2014) 117-123.
- [15] S. E. Fadugba, “Solution of Fractional Order Equations in the Domain of the Mellin Transform”, Journal of the Nigerian Society of Physical Sciences **4** (2019) 138. <https://doi.org/10.46481/jnsps.2019.31>
- [16] Y. Nawaz, “Variational iteration method and homotopy perturbation method for fourth-order fractional integro-differential equations”, Computers & Mathematics with Applications **8** (2011) 2330.
- [17] O. A. Uwaheren, A. F. Adebisi & O. A. Taiwo, “Perturbed Collocation Method For Solving Singular Multi-order Fractional Differential Equations of Lane-Emden Type”, Journal of the Nigerian Society of Physical Sciences **3** (2020) 141. <https://doi.org/10.46481/jnsps.2020.69>.
- [18] M. K. Shahooth, R. R. Ahmed, U-K. S. Din, W. Swidan, O. K. Al-Husseini & W. K. Shahooth, “Approximation Solution to Solving Linear Volterra-Fredholm Integro-Differential Equations of the Second Kind by Using Bernstein Polynomials Method”, J Appl Computat Math **5** (2016). <https://doi.org/10.4172/2168-9679.1000298>.
- [19] O. A. Uwaheren, A. F. Adebisi, O. T. Olotu, M. O. Etuk & O. J. Peter, “Legendre Galerkin Method for Solving Fractional Integro-Differential Equations of Fredholm Type”, The Aligarh Bulletin of Mathematics **40** (2021) 15.
- [20] T. Oyedepo, O. A. Uwaheren, E. P. Okperhie & O. J. Peter, “Solution of Fractional Integro-Differential Equation Using Modified Homotopy Perturbation Technique and Constructed Orthogonal Polynomials as Basis Functions”, Journal of Science Technology and Education **7** (2019). ISSN: 2277-0011; Journal homepage: www.atbuftejoste.com
- [21] O. A. Uwaheren, A. F. Adebisi, C. Y. Ishola, M. T. Raji, A. O. Yekeem & O. J. Peter, “Numerical Solution of Volterra Integro-differential Equation by Akbari-Ganji’s Method”, Journal of Mathematics and Its Application **16** (2022) 1123.
- [22] G. Ajileye, A. A. James, A. M. Ayinde & T. Oyedepo, “Collocation Approach for the Computational Solution Of Fredholm-Volterra Fractional Order of Integro-Differential Equations”, J. Nig. Soc. Phys. Sci. **4** (2022) 834.
- [23] G. Ajileye & F. A. Aminu, “Approximate Solution to First-Order Integro-differential Equations Using Polynomial Collocation Approach”, J Appl Computat Math. **11** (2022) 486.